

# The astrophysics of binary neutron star mergers

## Lecture III

Luciano Rezzolla

Institute for Theoretical Physics, Frankfurt



Zakopane  
17.06.19



# Plan of the lectures

\* Lecture I: the **math** of neutron-star mergers

\* Lecture II: the **physics** of neutron-star mergers


\* Lecture III: the **astrophysics** of neutron-star mergers

\* L. Baiotti and L. Rezzolla, Rep. Prog. Phys. 80, 096901, 2017

\* V. Paschalidis, Classical Quantum Gravity 34, 084002 2017


\* Rezzolla and Zanotti, *“Relativistic Hydrodynamics”*, Oxford University Press, 2013

# Electromagnetic counterparts



# Electromagnetic counterparts

- Since 70's we have observed flashes of gamma rays with enormous energies  $10^{50-53}$  erg: **gamma-ray bursts**.
- There are two families of bursts: “**long**” and “**short**”.
- The first ones last **tens** or more of **seconds** and could be due to the collapse of very massive stars.
- The second ones last **less** than a **second**.
- Merging neutron stars most reasonable explanation but how do you produce a **jet**?



# Electromagnetic counterparts (B-field)

**B-fields** essential for EMCs. Most simulations use **ideal MHD**: (infinite conductivity, B-field advected). Simple questions:

- can B-fields be measured during the inspiral?
- is EMC produced before merger?
- do B-fields grow after merger and yield EMC?
- does jet appear after BH formation and yield EMC?

Last two questions are **incredibly hard** to answer; may require far more sophisticated numerics and microphysics

# Electromagnetic counterpart (EMC)

B-fields essential for EMCs. Most simulations use **ideal MHD**: (infinite conductivity, B-field advected). Simple questions to ask:

- can B-fields be measured during the inspiral?



**NO!**

- is EMC produced before merger?



**Maybe. Luminosity is however low.**

- do B-fields grow after merger and yield EMC?



**Certainly but unclear how much:  $20-10^3$  amplification?**

- does jet appear after BH formation and yield EMC?



**YES (jet structure and outflow). Unclear how to produce ultrarelativistic outflow.**

Presence of a jet immediately implies presence of large-scale magnetic fields

What happens when magnetised stars collide?



Need to solve equations of magnetohydrodynamics in addition to the Einstein equations

$$T_{\mu\nu} = (e + p) u_{\mu} u_{\nu} + p g_{\mu\nu} + F_{\mu}^{\lambda} F_{\nu\lambda} - \frac{1}{4} g_{\mu\nu} F^{\lambda\alpha} F_{\lambda\alpha},$$

$$\nabla^{\nu} T_{\mu\nu} = 0$$

$$\nabla_{\nu} (F^{\mu\nu} + g^{\mu\nu} \psi) = I^{\mu} - \kappa n^{\mu} \psi, \quad \nabla_{\nu} (*F^{\mu\nu} + g^{\mu\nu} \phi) = -\kappa n^{\mu} \phi,$$

# Can we detect B-fields in the inspiral?



Compare B/no-B field:

- **inspiral** waveform is different but for unrealistic B-fields (i.e.  $B \sim 10^{17}$  G).

- **post-merger** waveform is different for all masses; strong B-fields delay the collapse to BH

Influence of B-fields on inspiral is **unlikely to be detected** for realistic fields



# Can we detect B-fields in the inspiral?

To quantify the differences and determine whether detectors will see a difference in the inspiral, we calculate the **overlap**

$$\mathcal{O}[h_{B1}, h_{B2}] \equiv \frac{\langle h_{B1} | h_{B2} \rangle}{\sqrt{\langle h_{B1} | h_{B1} \rangle \langle h_{B2} | h_{B2} \rangle}}$$

where the scalar product is


$$\langle h_{B1} | h_{B2} \rangle \equiv 4\Re \int_0^\infty df \frac{\tilde{h}_{B1}(f) \tilde{h}_{B2}^*(f)}{S_h(f)}$$

In essence, at these res:

$$\mathcal{O}[h_{B0}, h_B] \gtrsim 0.999$$

$$\text{for } B \lesssim 10^{17} \text{ G}$$

Influence of B-fields on inspiral is **unlikely to be detected**




Presence of a jet immediately implies presence  
of large-scale magnetic fields



What happens when magnetised stars collide?

Need to solve equations of  
magnetohydrodynamics in addition to the  
Einstein equations

If magnetic fields cannot be measured in the inspiral, what happens after merger?




$$M = 1.5 M_{\odot}, B_0 = 10^{12} \text{ G}$$




Animations: LR, Koppitz


# What happens when magnetised stars collide?



*Simulation begins*





*7.4 milliseconds*




*13.8 milliseconds*

Magnetic fields in the HMNS have complex topology: dipolar fields are destroyed.






Simulation begins



7.4 milliseconds



13.8 milliseconds



Black hole forms  
Mass: 2.9 suns  
Horizon diameter: 5.6 miles (9 km)



16.2 milliseconds







16.2 milliseconds

These simulations have shown that the merger of a magnetised binary has all the basic features behind SGRBs


$$J/M^2 = 0.83$$

$$M_{\text{tor}} = 0.063 M_{\odot}$$


$$t_{\text{accr}} \simeq M_{\text{tor}} / \dot{M} \simeq 0.3 \text{ s}$$




With due differences, other groups confirm this picture



Kiuchi+ 2014






Ruiz+ 2016



Kawamura+2016

Dionysopoulou+ 2015



RMHD



# Beyond IMHD: Resistive Magnetohydrodynamics

Dionysopoulou, Alic, LR (2015)

- Ideal MHD is a good approximation in the inspiral, but not after the merger; match to **electro-vacuum** not possible.
- Main difference in resistive regime is the current, which is dictated by Ohm's law but microphysics is **poorly** known.
- We know conductivity  $\sigma$  is a **tensor** but hardly know it as a scalar (prop. to density and inversely prop. to temperature).
- A simple prescription with scalar (isotropic) conductivity:

$$J^i = qv^i + W\sigma[E^i + \epsilon^{ijk}v_j B_k - (v_k E^k)v^i],$$



$\sigma \rightarrow \infty$  ideal-MHD (IMHD)

$\sigma \neq 0$  resistive-MHD (RMHD)

$\sigma \rightarrow 0$  electrovacuum







$$\sigma = f(\rho, \rho_{\min})$$



phenomenological prescription



$t = 19.892 \text{ ms}$

$t = 22.446 \text{ ms}$







NOTE: the **magnetic jet structure** is **not** an **outflow**. It's a plasma-confining structure.

In **IMHD** the magnetic jet structure is present but less regular.  
In **RMHD** it is more regular at all scales.


The magnetic jet structure maintains its coherence up to the largest scale of the system.

**RMHD**




# Ejected matter and nucleosynthesis

Bovard+ (2017)




# Nucleosynthesis


- Already in the 50's, nuclear physicists had tracked the production of elements in stars via nuclear fusion.
- **Heavy elements** ( $A \gtrsim 56$ ) cannot be produced in stellar interiors but can be synthesised during a **supernova**.
- SN simulations have shown that temperatures/energies not enough to produce “**very heavy**” elements ( $A \gtrsim 120$ ).
- To produce such elements very high temperatures and “**neutron-rich**” material is needed.
- **Neutron-star mergers** seem perfect candidates for this process!



log(density) (g/cm<sup>3</sup>)



$t = 11.801$  ms



L. Bovard, LR



# Ejection of mass

- After merger mass is lost in many different **channels** (shock heating, neutrino or magnetic-driven winds) and on very different **timescales** (**dynamical** and **secular**).

$$M_{\text{ej,blue}} \simeq 0.025 M_{\odot}$$
$$\beta_{\text{max,blue}} \lesssim 0.3$$

$$M_{\text{ej,red}} \ll 10^{-2} M_{\odot}$$
$$\beta_{\text{max,red}} \approx 0.1$$

$$M_{\text{ej,blue}} \approx 0$$

$$M_{\text{ej,red}} \gtrsim 10^{-2} M_{\odot}$$

DISK HMNS DISK

$$0.2 \lesssim Y_e \lesssim 0.3$$

$$Y_e \lesssim 0.2$$

before collapse

TORUS


TORUS

collimated jet  
cocoon

after collapse


# Relative abundances

- Mass ejection can either be **dynamical** (shocks; 100 ms) or **secular** (magnetic or neutrino-driven winds; 1-10 s).
- Even **tiny amounts** of ejected matter ( $0.01M_{\odot}$ ) sufficient to explain observed abundances.
- Abundances for  $A > 120$  good agreement with solar. **robust** for different **EOSs, masses, nuclear reactions and merger type**




# Relative abundances

- Mass ejection can either be **dynamical** (shocks; 100 ms) or **secular** (magnetic or neutrino-driven winds; 1-10 s).
- Even **tiny amounts** of ejected matter ( $0.01M_{\odot}$ ) sufficient to explain observed abundances.
- Abundances for  $A > 120$  good agreement with solar. **robust** for different **EOSs, masses, nuclear reactions and merger type**




# Relative abundances

- Mass ejection can either be **dynamical** (shocks; 100 ms) or **secular** (magnetic or neutrino-driven winds; 1-10 s).
- Even **tiny amounts** of ejected matter ( $0.01M_{\odot}$ ) sufficient to explain observed abundances.
- Abundances for  $A > 120$  good agreement with solar. **robust** for different **EOSs, masses, nuclear reactions and merger type**



- GW170817 produced total of **16,000** times the mass of the Earth in heavy elements (**10** Earth masses in **gold/platinum**)
- We are not only **stellar dust** but also **neutron-star dust!**


# Spatial distributions: $M_{ej}$ Bovard+ 17



Spatial distribution of  $M_{ej}$  impacts detectability of EM counterpart:

- ★ most of  $M_{ej}$  lost at low latitudes;
- ★ depending on EOS/mass, contamination also in polar regions

# Spatial distributions: $Y_e$ Bovard+ 17




Spatial distribution of  $Y_e$  impacts detectability of EM counterpart:


- ★ high  $Y_e$  in **polar** regions: **blue** (optical) macronova
- ★ low  $Y_e$  in **equatorial** regions: **red** (FIR) macronova

# Kilonova emission

- Ejected matter undergoes **nucleosynthesis** as expands and cools.
- When critical densities and temperatures are reached, matter undergoes radioactive decay emitting light (optical/infrared): **kilonova/macronova** (Li & Paczynski '98).



simulations




observations (Tanvir+2017)

- Astronomical observations of GW170817 show **kilonova emission**: evidence connection **GRBs** and **binary neutron stars!**

# When did the merger of GW170817 collapse to a BH?

Gill, Nathanail, LR (2019)






# Why is this important?

Conservative assumption: the remnant of GW170817 collapsed to a BH. GRB observed at  $t_{\text{del}} = 1.74 \pm 0.05 \text{ s}$

However, **when** did it actually collapse?

- If it collapsed **too early** it would have not **ejected** the matter that we can deduce from the kilonova emission.
- If it collapsed **too late** it would have not produced the delay we have observed of.
- The more the mass ejected, the longer for the jet to bore its way and **breakout**.



# Ejection of mass

- After merger mass is lost in many different **channels** (shock heating, neutrino or magnetic-driven winds) and on very different **timescales** (dynamical and secular).

$$M_{\text{ej,blue}} \simeq 0.025 M_{\odot}$$
$$\beta_{\text{max,blue}} \lesssim 0.3$$

$$M_{\text{ej,red}} \ll 10^{-2} M_{\odot}$$
$$\beta_{\text{max,red}} \approx 0.1$$

$$M_{\text{ej,blue}} \approx 0$$

$$M_{\text{ej,red}} \gtrsim 10^{-2} M_{\odot}$$

DISK HMNS DISK

$$0.2 \lesssim Y_e \lesssim 0.3$$

$$Y_e \lesssim 0.2$$

before collapse


collimated jet  
cocoon

TORUS

TORUS

after collapse

# Ejection of mass



- Shown are the mass-ejection rates deduced from numerical simulations.

- $M_{dyn}$ : matter ejected dynamically


- $M_{\nu}$  : matter ejected via neutrino-driven winds

- $M_B$ : matter ejected via magnetically driven winds

All channels have contribution from the central object and the disk.

All channels provide both **blue** or **red** ejecta in different amounts


# Constraints from mass ejection



- Shown are the mass contributions (blue/red) on “long” timescales.
- Blue ejecta essentially stops after collapse and constraints collapse time **from mass ejection** to be

$$t_{\text{coll}} = 1.14^{+0.60}_{-0.50} \text{ s}$$

# Constraints from breakout




- Breakout time depends on collapse time, speed of ejecta jet opening angle, and energy injected (more and faster ejecta, longer to escape).

- Given measured  $t_{\text{del}}$  we can constrain collapse time **from breakout** to be

$$t_{\text{coll}} = 0.82 \pm 0.15 \text{ s}$$

$$t_{\text{del}} = 1.74 \pm 0.05 \text{ s} = t_{\text{coll}} + t_{\text{br}}(t_{\text{coll}}) + t_R$$



# Putting things together

- Can combine two constraints and their uncertainties to obtain a single estimate

$$t_{\text{coll}} = 0.98^{+0.31}_{-0.26} \text{ s}$$


- What are the **implications?**

- \*correlates  $M_{\text{ej,blue}}$  and  $t_{\text{coll}}$ :  
to be tested new detections

- \*much longer than what can be simulated accurately ( $\sim 0.1$  s)

- \*mechanisms other than GWs for loss of angular momentum:  
spin down due to dipolar EM radiation appears reasonable

- \*this implies  $B \gtrsim 10^{16}$  G need to be produced **after** merger.



# Recap

- ✓ **Mergers** lead naturally to EM counterparts (GRB, kilonova).
- ✓ Magnetic fields unlikely to be detected during the inspiral but **important** after the merger: instabilities and EM counterparts.
- ✓ **Electromagnetic counterparts** and a **jet** are **likely** to be produced but the details of this picture are still **far from clear**.
- ✓ **Mergers** lead to tiny but important ejected matter and macronova emission.
- ✓ “high-*A*” nucleosynthesis very robust (little dependence on EOS and mass ratio) and good agreement with solar abundances.
- ✓ First constraints on lifetime of **GW170817** remnant

$$t_{\text{coll}} = 0.98^{+0.31}_{-0.26} \text{ s}$$